

Gene Section

Mini Review

RPS27 (ribosomal protein S27)

Tiffany Pierson, Brendan C Stack Jr

Department of Otolaryngology-Head and Neck Surgery, University of Arkansas for Medical Sciences, AR 72205, USA (TP, BCJrS)

Published in Atlas Database: August 2011

Online updated version : <http://AtlasGeneticsOncology.org/Genes/RPS27ID45550ch1q21.html>

DOI: 10.4267/2042/46937

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2012 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

Identity

Other names: MPS-1, MPS1, S27

HGNC (Hugo): RPS27

Location: 1q21.3

Local order: Human RPS27 is found on chromosome 1: 153963235 - 153964626 bp from pter. Information about the local order for RPS27 can be found at ensembl.org. Four transcripts have been identified, but only the first will be discussed below.

DNA/RNA

Description

The RPS27 gene is comprised of 1.39 kb and consists of 4 exons. This gene is a member of the Human CCDS set: CCDS1059.

Transcription

The transcript is 350 base pairs long.

Pseudogene

Multiple RPS27 pseudogenes are dispersed throughout the genome. The RPS27L pseudogene, located at 15q22.2, is known to encode a protein that shares 96% of its amino acid sequence with RPS27 (Balasubramanian et al., 2009).

Protein

Description

RPS27 is a 9461 Da protein composed of 84 amino acids. The protein contains a C4 zinc finger

domain, similar to steroid and thyroid hormones, which enables DNA binding. RPS27 is found in both the cytoplasm and the nucleus.

Expression

Ubiquitous expression. Expressed at high levels in actively dividing cells and in cancers of ectodermal origin, as well as in melanoma (Fernandez-Pol et al., 1993). When overexpressed, it is secreted into serum (Lee et al., 2004).

Function

1. Component of the 40S ribosomal subunit in the cytoplasm: ribosomes carry out translation of proteins. The eukaryotic ribosome is made up of a small 40S and a large 60S subunit. Together these subunits are comprised of 4 different rRNA species and almost 80 different RP's (ribosomal proteins). As a component of the 40S subunit, RPS27 is found near RPS18 and covalently bound to translation initiation factor eIF3.
2. A mediator of cellular proliferation and survival: expression is induced by a variety of growth factors and other signaling molecules, including TGF-beta and cAMP; RPS27 can bind to cAMP response elements of DNA (Fernandez-Pol et al., 1993).
3. Oncogenesis (see below).

Homology

Member of the ribosomal protein S27e family.

Mutations

Note

Single nucleotide polymorphisms have been identified, but have not been linked to disease.

Implicated in

Various carcinomas and melanoma

Note

RPS27 overexpression has been reported in many cancers including prostate cancer (Fernandez-Pol et al., 1997), colorectal cancer (Ganger et al., 1997), liver cancer (Ganger et al., 2001), breast cancer (Atsuta et al., 2002), head and neck squamous cell cancer (HNSCC) (Stack et al., 1999; Stack et al., 2004; Lee et al., 2004), gastric cancer (Wang, et al., 2006), as well as, melanoma (Santa Cruz et al., 1997).

Since high serum levels of RPS27 have been found in cancer patients, especially in head and neck squamous cell carcinoma (HNSCC), the protein can be used as a tumor marker (Fernandez-Pol et al., 1996; Lee et al., 2004; Stack et al. 2004).

Prognosis

It was reported that RPS27 levels correlate with tumor stage in patients with gastric cancer, thus high levels serve as a poor prognostic indicator (Wang et al., 2006).

Oncogenesis

The mechanism behind RPS27 overexpression is currently under investigation. One explanation recently offered arises from the relationship between RPS27, MDM2 and p53: RPS27 is a p53 repressible protein (He and Sun, 2007; Li et al., 2007). A 2011 study found that it competes with p53 for a central acidic binding domain on MDM2. Once bound, MDM2 is stimulated to ubiquitinate and degrade the RPS27 or p53, whichever it is bound to. When RPS27 levels are elevated, it can out-compete p53 for MDM2 binding and subsequent degradation, thus stabilizing p53 levels. This would be an appropriate cellular response to genotoxic stress. The same study also found that mutant p53 cannot suppress RPS27, only the wild-type can. Since mutated p53 is found in almost 50% of all human cancers, RPS27 overexpression logically follows. Furthermore, stabilization of mutant p53 levels associated with RPS27 abundance could provide malignant cells with a growth advantage (Xiong et al., 2011).

RPS27 knockdown was found to enhance spontaneous apoptosis of tumor cells via caspase-3 activation (Wang et al., 2006; Yang et al., 2011).

HNSCC: some have questioned if RPS27 overexpression is the cause or result of cancer. A 2010 study overexpressed RPS27 in a line of HNSCC cells to study the impact on tumor behavior. They found that RPS27 overexpression resulted in reduced cancer cell growth, proliferation rate and angiogenesis. RPS27 overexpression was also found to reduce the mRNA of Paxillin, a focal adhesion protein up regulated in HNSCC and many other cancer cells. RPS27 induced Paxillin repression offers a possible explanation for the decreased HNSCC growth (Dai et al., 2010).

References

- Fernandez-Pol JA, Klos DJ, Hamilton PD. A growth factor-inducible gene encodes a novel nuclear protein with zinc finger structure. *J Biol Chem*. 1993 Oct 5;268(28):21198-204
- Fernandez-Pol JA, Fletcher JW, Hamilton PD, Klos DJ. Expression of metallopanstimulin and oncogenesis in human prostatic carcinoma. *Anticancer Res*. 1997 May-Jun;17(3A):1519-30
- Ganger DR, Hamilton PD, Fletcher JW, Fernandez-Pol JA. Metallopanstimulin is overexpressed in a patient with colonic carcinoma. *Anticancer Res*. 1997 May-Jun;17(3C):1993-9
- Santa Cruz DJ, Hamilton PD, Klos DJ, Fernandez-Pol JA. Differential expression of metallopanstimulin/S27 ribosomal protein in melanocytic lesions of the skin. *J Cutan Pathol*. 1997 Oct;24(9):533-42
- Stack BC Jr, Dalsaso TA, Lee C Jr, Lowe VJ, Hamilton PD, Fletcher JW, Fernandez-Pol JA. Overexpression of MPS antigens by squamous cell carcinomas of the head and neck: immunohistochemical and serological correlation with FDG positron emission tomography. *Anticancer Res*. 1999 Nov-Dec;19(6C):5503-10
- Ganger DR, Hamilton PD, Klos DJ, Jakate S, McChesney L, Fernandez-Pol JA. Differential expression of metallopanstimulin/S27 ribosomal protein in hepatic regeneration and neoplasia. *Cancer Detect Prev*. 2001;25(3):231-6
- Atsuta Y, Aoki N, Sato K, Oikawa K, Nochi H, Miyokawa N, Hirata S, Kimura S, Sasajima T, Katagiri M. Identification of metallopanstimulin-1 as a member of a tumor associated antigen in patients with breast cancer. *Cancer Lett*. 2002 Aug 8;182(1):101-7
- Lee WJ, Keefer K, Hollenbeak CS, Stack BC Jr. A new assay to screen for head and neck squamous cell carcinoma using the tumor marker metallopanstimulin. *Otolaryngol Head Neck Surg*. 2004 Oct;131(4):466-71
- Stack BC Jr, Hollenbeak CS, Lee CM, Dunphy FR, Lowe VJ, Hamilton PD. Metallopanstimulin as a marker for head and neck cancer. *World J Surg Oncol*. 2004 Dec 14;2:45
- Gilkes DM, Chen L, Chen J. MDMX regulation of p53 response to ribosomal stress. *EMBO J*. 2006 Nov 29;25(23):5614-25
- Wang YW, Qu Y, Li JF, Chen XH, Liu BY, Gu QL, Zhu ZG. In vitro and in vivo evidence of metallopanstimulin-1 in gastric cancer progression and tumorigenicity. *Clin Cancer Res*. 2006 Aug 15;12(16):4965-73
- He H, Sun Y. Ribosomal protein S27L is a direct p53 target that regulates apoptosis. *Oncogene*. 2007 Apr 26;26(19):2707-16
- Li J, Tan J, Zhuang L, Banerjee B, Yang X, Chau JF, Lee PL, Hande MP, Li B, Yu Q. Ribosomal protein S27-like, a p53-inducible modulator of cell fate in response to genotoxic stress. *Cancer Res*. 2007 Dec 1;67(23):11317-26
- Balasubramanian S, Zheng D, Liu YJ, Fang G, Frankish A, Carriero N, Robilotto R, Cayting P, Gerstein M. Comparative analysis of processed ribosomal protein pseudogenes in four mammalian genomes. *Genome Biol*. 2009;10(1):R2
- Dai Y, Pierson SE, Dudney WC, Stack BC Jr. Extraribosomal function of metallopanstimulin-1: reducing paxillin in head and neck squamous cell carcinoma and inhibiting tumor growth. *Int J Cancer*. 2010 Feb 1;126(3):611-9
- Xiong X, Zhao Y, He H, Sun Y. Ribosomal protein S27-like and S27 interplay with p53-MDM2 axis as a target, a substrate and a regulator. *Oncogene*. 2011 Apr 14;30(15):1798-811

Yang ZY, Qu Y, Zhang Q, Wei M, Liu CX, Chen XH, Yan M, Zhu ZG, Liu BY, Chen GQ, Wu YL, Gu QL. Knockdown of metalloproteinase-1 inhibits NF- κ B signaling at different levels: The role of apoptosis induction of gastric cancer cells. *Int J Cancer*. 2011 Jul 27; doi: 10.1002/ijc.26331

This article should be referenced as such:

Pierson T, Stack BCJr. RPS27 (ribosomal protein S27). *Atlas Genet Cytogenet Oncol Haematol*. 2012; 16(2):103-105.
